arXiv:cond-mat/0702344v2 [cond-mat.supr-con] 5 Jun 2007

Half quantum vortex in superfluid ^{3}He-A phase in parallel plate geometry

Hae-Young Kee[1] and Kazumi Maki[2]
[1] Department of Physics, University of Toronto, Toronto, Ontario M5S 1A7, Canada
[2] Department of Physics and Astronomy, University of Southern California, Los Angeles, CA 90089-0484, USA

The half quantum vortex(HQV) in condensate has been studied, since it was predicted by Salomaa and Volovik in superfluid ^{3}He-A phase. However, an experimental evidence for its existence has not been reported so far. Motivated by a recent experimental report by Yamashita et al[1], we study the HQVs in superfluid ^{3}He confined between two parallel plates with a gap D $\sim$ 10 μm in the presence of a magnetic field H $\sim$ 26 mT perpendicular to the parallel plates. We find that the bound HQVs are more stable than the singular vortices and free pairs of HQVs, when the rotation perpendicular to the parallel plates is below the critical speed, $\Omega_c \sim 2$ rad/s. The bound pair of HQVs accompanies the tilting of $\hat{d}$-vector out of the plane, which leads to an additional absorption in NMR spectra. Our study appears to describe the temperature and rotation dependence of the observed satellite NMR signal, which supports the existence of the HQVs in ^{3}He.

PACS numbers: 67.57.-z,67.57.Fg,67.57.Lm

I. INTRODUCTION

The superfluid ^{3}He-A is a remarkable condensate with several broken symmetries and a variety of topological defects. It was proposed that many concepts and notions generated in the high energy particle physics can be tested in the earth bound ultra-low temperature laboratories.[2,3,4]. Indeed a variety of vortices, which include the continuous vortices (the vortices with soft-core) have been observed in the rotating superfluid ^{3}He-A.[5,6] All of the continuous vortices are associated with $\hat{l}$-texture are stable in the open systems where angular momentum of Cooper pair, $\hat{l}$-vector is not constrained.

On the other hand, in the parallel plate geometry, where superfluid ^{3}He is confined between two parallel plates with a gap $D \sim \xi_D$, where ξ_D (~ 10 μm) is the dipole coherence length, the $\hat{l}$-vector is forced to be perpendicular to the plate.[7,9] In such a geometry, the continuous vortices cannot exist, and one may conclude that the vortices with the circulation $N = \pm 1$ called the phase vortices or the singular vortices are only possible vortices.[2] However, Salomaa and Volovik[10] predicted that the half quantum vortices(HQVs) with $N = \pm\frac{1}{2}$ should exist in the parallel plate geometry. In spite of intensive experimental search, the HQV has not been found in the rotating superfluid ^{3}He-A in the confined geometry.[6]

It is worthwhile to note that the HQVs have played an important role outside the superfluid ^{3}He-A. For example, the HQVs in the tricrystal geometry in the high temperature cuprate superconductors played the crucial role in identifying d-wave symmetry of the high temperature superconducting order parameter.[11] The HQVs in triplet superconductors have been also suggested[12,13,14] to interpret the extremely strong vortex pinning observed in triplet superconductors such as UPt$_3$, U$_{1-x}$Th$_x$Be$_{13}$, and Sr$_2$RuO$_4$.[15]

The present work is motivated by the recent experiment by Yamashita et al.[1] We consider the superfluid ^{3}He-A confined between the two parallel plate with the gap $D \sim \xi_D$, where the rotation axis and the magnetic field are perpendicular to the parallel plates. Using the standard formula for free energies given in [3,13], we found the followings. (a) The singular vortices with $N = \pm 1$ are unstable and split into pairs of HQV with $N = \pm\frac{1}{2}$, unless the rotation speed is very high, $\Omega > \Omega_u \sim 5 \times 10^6$ rad/s. (b) For small rotation speed ($\Omega \leq \Omega_c \sim 2$ rad/s), the bound pairs of HQVs are more stable than the free HQVs. (c) Only the bound pairs of HQVs are detectable by NMR as satellite signals. Both the singular vortices and free HQVs lead to NMR signals indistinguishable from backgrounds, since they do not disturb the configuration of $\hat{l} \perp \hat{d}$. (d) Our results of the satellite frequencies and the intensities are consistent with the observation of satellite peak in the parallel plate geometry.[1] In particular, while Ω_c weakly depends on the temperature and the magnetic field strength, both the satellite frequency and the satellite intensity are almost independent of the field strength and temperature.

II. FREE ENERGIES OF ONE SINGULAR VORTEX AND A PAIR OF HQV

The superconducting order parameter of superfluid ^{3}He-A is characterized by 2 unit vectors called l-vector, $\hat{l}$ and d-vector, $\hat{d}$. We can write the gap function as[3,4]

$$\Delta_{ab}(\mathbf{k}) = \vec{D}(\mathbf{k}) \cdot (\vec{\sigma} i \sigma_2)_{ab} \tag{1}$$

where

$$\vec{D}(\mathbf{k}) = \Delta \hat{d}(\hat{k}_1 + i\hat{k}_2) \tag{2}$$

Here σ_μ ($\mu = 1,2,3$) are Pauli matrices and a, b are spin ↑ and ↓. $\hat{k}_i$ ($i = 1,2$) are the projection of the unit wave vector $\hat{k}$ in the plane perpendicular to $\hat{l}$. Therefore $\hat{l}$ corresponds to the quantization axis of the angular momentum of the Cooper pair. This is also called the chiral vector.

When the spatial variation of $\hat{d}$ and $\hat{l}$ are much slower than ξ ($\sim 0.01\mu m$), coherence length, the spatial configuration of $\hat{l}$ and $\hat{d}$ are described by the texture free energy.[3]. In the following, we shall compare the texture free energy of a singular vortex(or phase vortex), and a free pair of HQV with an optimized distance R, where R is a distance between two HQVs.

The texture free energy is given by

$$F = \frac{1}{2}\chi_N C^2 \int dxdy \left(K(\nabla\Phi)^2 + \sum_{ij} |\partial_i \hat{d}_j|^2 + \xi_H^{-2} \hat{d}_z^2 \right), \tag{3}$$

where we assumed the parallel plates extend in the $x-y$ plane. Here χ_N and C are spin susceptibility and spin wave velocity. Then K is given by the ratio between the superfluid density, ρ_s and spin superfluid density, ρ_{sp}.

$$K = \frac{\rho_s}{\rho_{sp}} = \frac{1+\frac{1}{3}F_1}{1+\frac{1}{3}F_1^a} \frac{1+\frac{1}{3}F_1^a(1-\rho_s^0)}{1+\frac{1}{3}F_1(1-\rho_s^0)} \tag{4}$$

where F_1 and F_1^a are Landau's Fermi liquid coefficients. We also assumed $\hat{l} \parallel \hat{z}$ everywhere due to the parallel plate constraint.[7] Within the weak-coupling theory of BCS p-wave superconductor, we obtain

$$\rho_s^0 = \begin{cases} 2(1-t) & \text{for} \quad t \sim 1 \\ 1 - O(t^3) & \text{for} \quad t \to 0 \end{cases} \tag{5}$$

where $t = T/T_c$. For the pressure of 27.3 bar with the superfluid transition temperature $T_c = 2.46$ mK corresponding to the experimental set-up by Yamashita et al[1], we found

$$K = 1 + 2.295(1-t), \tag{6}$$

where $(1-t) << 1$.

When the parallel plates are rotated with the rotation axis perpendicular to the plates and rotation speed Ω, the density of $N = 1$ vortices is given by

$$n_v = (\pi a^2)^{-1} = 2\Omega/(p\kappa) \tag{7}$$

where $\kappa = 2\pi\hbar/(2m_{^3He}) = 6.59 \times 10^{-8}\mathrm{m}^2/\mathrm{s}$ is the quantum of circulation, and p is the number of quanta per vortex. $2a$ corresponds roughly the distance between two vortices, if we divide the two-dimensional space in circular cells with radius a. Therefore a is related to Ω by $a = \sqrt{p\kappa/(2\pi\Omega)}$. The free energy for one singular vortex in unit cell is given by

$$F_s = \pi \chi_N C^2 K \ln\left(\frac{a}{\xi}\right), \tag{8}$$

which does not involve the texture energy except $\Phi = \tan^{-1}(y/x)$. Here $\xi \sim 0.01$ μm is the coherence length.



FIG. 1: Phase diagram for SV and free HQV phases as functions of K and a/ξ. Singular vortex (SV) phase is stable only for small a/ξ for all temperatures.

On the other hand, the free energy for free pairs of HQVs with their distance $R(< 2a)$ is obtained as follows.

$$F_{fH} = \frac{\pi}{2}\chi_N C^2 \left\{ K \ln\left(\frac{a + \sqrt{a^2 - R^2/4}}{2\xi} \right) + \ln(R/\xi) - \frac{R}{4a} \arcsin\left(\frac{R}{\sqrt{a^2 + R^2/4}} \right) \right\} \tag{9}$$

To get the above free energy, we used the following forms for Φ and $\hat{d}$.

$$\Phi = \frac{1}{2}\left[\tan^{-1}\left(\frac{x+R/2}{y}\right) + \tan^{-1}\left(\frac{x-R/2}{y}\right) \right] \tag{10}$$

and

$$\hat{d} = \hat{x}\cos\alpha + \hat{y}\sin\alpha \tag{11}$$

with

$$\alpha = \frac{1}{2}\left[\tan^{-1}\left(\frac{x+R/2}{y}\right) - \tan^{-1}\left(\frac{x-R/2}{y}\right) \right] \tag{12}$$

As in the case of the singular vortex, this solution has $\hat{d}_z = 0$. If we neglect the last term in the Eq. (9), F_{fH} can be easily optimized for $R/(2a) = \sqrt{2K+1}/(K+1)$ and

$$F_{fH} \approx \frac{\pi}{2}\chi_N C^2 \left\{ K \ln\left(\frac{a}{2\xi}\frac{1+2K}{1+K} \right) + \ln\left(\frac{2a}{\xi}\frac{\sqrt{1+2K}}{2(1+K)} \right) - \frac{2\sqrt{1+2K}}{1+K} \arcsin\left(\frac{\sqrt{1+2K}}{\sqrt{2+4K+K^2}} \right) \right\}. \tag{13}$$

Now we compare the free energies of singular vortex, Eq (8) and pairs of free HQV, Eq. (13), and find that the free HQVs are stable than the singular vortices if a/ξ satisfies the following equation.

$$\frac{a}{\xi} \geq \frac{1+2K}{2(1+K)} \left(\frac{(1+2K)^{3/2}}{(1+K)^2} \right)^{\frac{1}{(1-K)}} \tag{14}$$

In the above equation, the last term in Eq. (13) is neglected for the analytic solution.[8] However, we find that



FIG. 2: A configuration of $\hat{d}$-vector for $\beta = 0$

this approximation does not make a qualitative difference for the phase diagram as a function of (a/ξ) and K shown in Fig 1. The free HQV is stable than the singular vortex for all temperatures ($1 < K < 3.295$), since we are interested in small rotations of $\Omega << \Omega_u$, where $\Omega_u \sim 5 \times 10^6$ rad/s which corresponds to $a \sim \xi \sim 0.01\mu$m. Hereafter, we will limit ourselves to HQVs and compare the free energies between free HQVs and bound HQVs.

III. FREE ENERGY OF THE BOUND HQVS PAIRS

If we allow $\hat{d}$-vector to bend out of the $x - y$ plane in the vicinity of $\hat{d}$-digyrations, the system will lower the energy. So it is quite possible that two HQVs are bound when they are close enough. Let us consider the following $\hat{d}$ configurations.

$$\hat{d} = -\cos\alpha(\hat{x}\cos\beta + \hat{y}\sin\beta) + \sin\alpha\hat{z} \tag{15}$$

where β is the cyclic variable, and an example of $\hat{d}$ configurations is shown in Fig. 2 for $\beta = 0$.

Using the free energy formula for the bound HQVs, Eq. (8) in [13], we find

$$\begin{aligned} F_{bH} &= \frac{\pi}{2}\chi_N C^2 \left\{ K\ln\left(\frac{a + \sqrt{a^2 - R^2/4}}{2\xi}\right) + \ln\left(\frac{R}{\xi}\right) \right. \\ &\left. -\frac{R}{4a}\arcsin\left(\frac{R}{\sqrt{a^2 + R^2/4}}\right) + \left(\frac{R}{2\xi_H}\right)^2 \ln\left(\frac{4\xi_H}{R}\right) \right\} \end{aligned} \tag{16}$$

The free energy can be optimized for R/a for given K and a/ξ_H assuming that $a/\xi_H >> 1$. We found that $R/a = 0.3$ for $a/\xi_H = 10$ for all K values, and $R/a = 0.15$ for $a/\xi_H = 20$ for all K values. For $a/\xi_H = 30$, $R/a = 0.1$ for all K values. The ratio, $R/\xi_H (= R/a \times a/\xi_H) \sim 3.0$ is independent of K and a/ξ_H. It is remarkable that R/ξ_H is independent of temperatures.

Using the optimal value of R/ξ_H, the free energy of the bound HQV can be written as

$$\begin{aligned} F_{bH} &= \frac{\pi}{2}\chi_N C^2 \left\{ K\ln\left(\frac{a}{\xi}\right) + \ln\left(3\frac{\xi_H}{\xi}\right) \right. \\ &\left. -\frac{3}{4}\frac{\xi_H}{a}\arcsin\left(\frac{3\xi_H}{a\sqrt{1 + \frac{9\xi_H^2}{4a^2}}}\right) + \frac{9}{4}\ln\left(\frac{4}{3}\right) \right\} \end{aligned} \tag{17}$$



FIG. 3: Phase diagram for bound HQV phase, and free HQV phase as functions of K and a/ξ_H. Note that the free HQV is only stable for a relatively small a/ξ_H for all K

We compare the free energies of bound HQV, Eq. (17) and free HQV, Eq. (13) neglecting the correction terms of $\hat{d}$ textures (the terms with arcsin(..)). We find that the bound HQVs is stable over the free HQV, if a/ξ_H satisfies the following equation.

$$\frac{a}{\xi_H} \geq \frac{8}{3}(2)^K(1 + K)^{1+K}(1 + 2K)^{-(K+\frac{1}{2})} \tag{18}$$

The phase diagram for free HQV and the bound HQV phases as a function of a/ξ_H and K is shown in Fig. 3. We see that the bound HQV is more stable than the free HQV when $a \geq 10\xi_H$ for all temperatures. In other words, the bound pair HQVs are always more stable than free HQVs for relatively small rotation, $\Omega < \Omega_c$, where $\Omega_c \sim 2$ rad/s, assuming that $\xi_H \sim 5$ μm. From these results, we analyze the Ω dependence of the satellite intensity (I_g) as follows.

In the small rotation speed, the bound pairs are most stable. So in the equilibrium condition, the satellite intensity increases linearly with Ω. When Ω reaches Ω_c, the free HQV pairs become more stable. Many bound pairs are converted into free pairs. This looks a case of phase separation. In order to clarify this situation, NMR experiment with the field gradient in the parallel plates is highly desirable. It is also interesting to note that the Ω dependence of the satellite intensity I_g looks surprisingly similar to $-4\pi M$ versus B in typical type II superconductors as first formulated by Abrikosov.[17]

IV. SPIN WAVE SPECTRUM

As we mentioned above, both the singular vortices and free HQVs do not produce any characteristic NMR signals, since their presence does not disturb the configuration of $\hat{l} \perp \hat{d}$. On the other hand, the bound pairs of HQVs are clearly visible by NMR as discussed in [13,16]. We adopt the formulation for the NMR satellite frequencies and intensities studied in [16]. However, in the present case there will be no potential energy for the longitudinal mode, since the rotation of $\hat{d}$ around the z-axis does not change the dipole energy. In other words, the bound pairs of HQVs as well as singular vortices and free HQVs are invisible by longitudinal NMR. All of them give the no longitudinal resonance. In contrast, the transverse NMR is of great interest.

The transverse resonance frequency, λ_g is determined as follows. Comparing to [16], we consider the bound HQVs when $\vec{H} \parallel \hat{z}$ and $\xi_H < \xi_D$. Let us consider a small oscillation of $\hat{d}$ as follows.

$$\begin{aligned}\hat{d} + \delta\hat{d} \;&=\; \cos(\alpha + g)\left[\cos(\beta + f)\hat{x} + \sin(\beta + f)\hat{y}\right] \\ &\quad + \sin(\alpha + g)\hat{z}. \qquad (19)\end{aligned}$$

The change of free energy due to the small oscillation up to the quadratic order in f and g is found as follows.

$$\begin{aligned}\delta F = \sum_{i,j} |\nabla_i \delta\hat{d}_j|^2 + \xi_H^{-2}|\delta d_z|^2 \;&=\; \cos^2(\alpha)|\vec{\nabla} f|^2 + |\vec{\nabla} g|^2 \\ &+\; \xi_H^{-2}(1 - 2\sin^2(\alpha))g^2. \qquad (20)\end{aligned}$$

Therefore, the satellite frequencies of λ_f and λ_g are obtained by the following eigenvalue equations.

$$\begin{aligned}-\,\xi_H^2 \nabla(\cos^2(\alpha)\nabla f) \;&=\; \lambda_f \cos^2(\alpha) f \\ -\xi_H^2 \nabla^2 g - (1 - 2\sin^2(\alpha))g \;&=\; \lambda_g g, \qquad (21)\end{aligned}$$

As we mentioned above, the longitudinal mode has a trivial solution of $f = \textit{constant}$ with $\lambda_f = 0$. Therefore, there will be no longitudinal resonance. As to the transverse mode, the potential $V_g(x, y) = -1 + 2\sin^2(\alpha)$ is shown in Fig. 4, and $\sin^2(\alpha)$ is given by

$$\sin^2(\alpha) = \frac{\sin^2 v}{\sinh^2 u + \sin^2 v} \qquad (22)$$

using the change of parameters as

$$x = \frac{R}{2}\cosh u \cos v, \qquad y = \frac{R}{2}\sinh u \sin v, \qquad (23)$$

Note that the potential energy in the present case is completely different from the one considered in [16]. This implies (a) first of all spin waves are extended, and (b) the related intensities are $\sim \pi a^2 n_v$. At the critical rotation of $\Omega = \Omega_c$, $n_v = (\pi a^2)^{-1}$.



FIG. 4: Potential for the transverse mode is shown in $x - y$ plane. It is repulsive between two HQVs, where R is the optimal separation between the bound HQVs.



FIG. 5: The satellite frequency of the transverse mode as a function of ξ_H/R is shown for the bound HQVs.

These eigenequations can be solved using a variational function. Note that there is a trivial solution of $g_0 = \mathrm{sech}u$ with $\lambda_g = -1$, which does not give a distint satellite signal. So we look for a solution which is orthogonal to g_0 and have a node in the spin wave function. We find the following function of g with a vartional parameter μ satisfies the above criteria.

$$g = (\mathrm{sech}u)^{\nu} - p \times \tanh u \times (\mathrm{sech}u)^{\mu}, \qquad (24)$$

where $\nu = 1.13$ and $p = (\mu^2 - 1)(B[\frac{1}{2}, \frac{\nu-1}{2}] - \frac{1}{2}B[\frac{1}{2}, \frac{\nu+1}{2}])/(\mu + 3)$, and B is the Euler beta function. With the above variational functions, we find the optimal value of λ_g which is obtained by minimizing λ_g with respect to μ. λ_g vs. ξ_H/R is shown in Fig. 5. Note that $R/\xi_H \sim 3$ is independent of a/ξ_H and K, as long as the system is in the regime where the bound HQVs are stable. The value of $\lambda_g \sim -0.05$ for $R/\xi_H = 3$ is close to the satellite frequency, $\lambda_g = -0.06 \sim -0.09$, observed in [1]. On the other hand, we find that the bound state does not exist, when $\xi_H/R > 0.35$.

The intensity of the transvermode is obtained as

$$I_g = \frac{n_v}{2}\frac{|\int_0^{arccosh(2a/R)} e^{i\alpha} g du|^2}{\int_0^{arccosh(2a/R)} |g|^2 du}. \qquad (25)$$

For $R/\xi_H = 3$, I_g depends on a/ξ_H. We find that $I_g = 23.4(R/2)^2 n_v$ and $I_g = 43.2(R/2)^2 n_v$ for $a/\xi_H = 7$, and $a/\xi_H = 10$, respectively. Since $n_v = (\pi a^2)^{-1}$ at $\Omega = \Omega_c$, I_g at the critical rotation is given by

$$\begin{aligned} I_g \;&=\; \frac{43.2}{\pi}\left(\frac{3}{20}\right)^2 = 0.31 \quad \text{for} \quad a/\xi_H = 10, \\ &=\; \frac{23.4}{\pi}\left(\frac{3}{14}\right)^2 = 0.34 \quad \text{for} \quad a/\xi_H = 7. \qquad (26)\end{aligned}$$

The ratio between the satellite intensity and total intensity, $\delta I/I$ in the NMR spectra was observed as $\delta I/I = 0.2$

at $T = 0.8T_c$, and 0.1 at $T = 0.93T_c$.[1] Therefore, the observed signal is about 60 % and 30 % of the theoretical value at $T = 0.8T_c$ and $T = 0.93T_c$, respectively. In our analysis, I_g does not explictily depend on temperature, while it is possible that the vortex pinning changes the satellite intensity. Therefore, the temperature dependence in the experiment suggests that the vortex pinning becomes weaker as temperature decreses.

V. DISCUSSION AND SUMMARY

Since Salomaa-Volovik's proposal for the HQVs in 1985, HQVs have been searched in the rotating superfluid ^{3}He-A in the confined geometry. However, all of earlier experiments have been done in the wider gap D ($> 25\mu m$). Then the most common texture is the vortex sheet associated with the composite soliton.[19] In the perspective, the vortices seen in Walmsley et al[18] appears to be the vortex sheet as well. The experimental data in [1] is quite different from the earlier ones in the following ways. (a) A new transverse satellite with $\lambda_g = -0.06 \sim -0.09$ is observed. (b) Both the satellite frequency and the critical value of Ω_c where the maximum value of $\delta I/I$ is observed are almost independent of temperature. (c) The critical value of Ω_1 where the first appearance of the satellite singal occurs and the maximal value of $\delta I/I$ appear to weakly depend on temperature.

The above observations are in fact consistent with our HQV scenario. We have shown that $R/\xi_H \sim 3.0$ is independent of H and T. Also from the universality of this ratio, we predict that the satellite frequency and the intensity are almost independent of H and T, when we ignore the vortex pinning effect. On the other hand, any texture involving $\hat{l}$-texture should depend strongly on temperature and gap size D. Furthermore, another satellite frequencies associated with $\hat{l}$-bend textures should appear near $\lambda_g \sim -1$.[9] Absence of these satellites clearly indicate that the satellites seen in Yamashita et al[1] should be due to the bound pairs of HQVs. The parallel experiments with different size of D are highly desirable to distinguish the $\hat{d}$-texture vs. $\hat{l}$-textures as well, as our study indicates that the satellite frequency is independent of D.

Acknowledgments

We would like to thank Takao Mizusaki and Minoru Yamashita for providing us their data on the ongoing superfluid ^{3}He experiments at ISSP, University of Tokyo at Kashiwanoha and their useful correspondences, which motivated the present study. This work was supported by NSERC of Canada, Canada Research Chair, Canadian Institute for Advancded Research, and Alfred P. Sloan Research Fellowship(HYK). One of us (KM) thanks Hyekyung Won at Hallym University for hospitality, where a part of the work was carried out.

---

[1] M. Yamashita, K. Izumina, A. Matsubara, Y. Sasaki, O. Ishikawa, T. Takagi, M. Kubota, and T. Mizusaki, LT24 Proceedings; AIP Proceedings **850**, 185 (2006); Private communication (M. Yamashita and T. Mizusaki).

[2] M. M. Salomaa and G. E. Volovik, Rev. Mod. Phys. **59**, 533 (1987).

[3] D. Vollhardt and P. Wölfle, The superfluid Phase of Helium 3 (Taylor and Francis, New York 1990); A. Leggett, Rev. Mod. Phys. **47**, 331 (1975).

[4] G. E. Volovik, Exotic Properties of Superfluid Helium 3 (World Scientific, Sigapore 1991).

[5] P. Hakonen, O.V. Lounasmaa and J. Simola, Physica B **160**, 1 (1989).

[6] G. E. Volovik, J. Low Temp. Phys. **121**, 357 (2000).

[7] V. Ambegaokar, P. G. de Gennes and D. Rainer, Phys. Rev. A **9**, 2676 (1974).

[8] Note when K approaches to 1, a/ξ approaches to 0, since $1/(1-K)$ becomes $-\infty$

[9] R. Bruinsma and K. Maki, J. Low Temp. Phys. **34**, 344 (1979).

[10] M. M. Salomaa and G. E. Volovik, Phys. Rev. Lett. **55**, 1184 (1985).

[11] C. C. Tsuei and J. R. Kirtley, Rev. Mod. Phys. **72**, 969 (2000).

[12] M. Sigrist and D. E. Agterberg, Prog. Theor. Phys. (Kyoto), **101**, 965 (1999).

[13] H. Y. Kee, Y. B. Kim, and K. Maki, Phys. Rev. B **62**, R9275 (2000).

[14] K. Maki, S. Haas, D. Darker, and H. Won in "Topology in Ordered Phases" edited by S. Tanda et al (World Scientific, Sigapore 2006).

[15] A. Amann and A. C. Mota, Phys. Rev. B **57**, 3640 (1998); E. Dumont and A. C. Mota, Phys. Rev. B **65**, 144519 (2000).

[16] C. R. Hu and K. Maki, Phys. Rev. B **36**, 6871 (1987).

[17] A. A. Abrikosov, Soviet Phys. JETP **5**, 1174 (1957).

[18] P. M. Walmsley, I. J. White and A. I. Golov, Phys. Rev. Lett. **93**, 195301 (2004).

[19] M. T. Heinilä and G. E. Volovik, Physica B **210**, 300 (1995).